

-continued

Pro	Glu	Val	Leu	Val	Gln	Glu	Val	Ile	Asp	Gln	Leu	Lys	Ala	Trp	Gly
	275						280					285			
Gly	Glu	Thr	Thr	Ser	Val	Arg	Glu	Asn	Ser	Gly	Ile	Glu	Glu	Lys	Val
	290					295					300				
Val	Phe	Ser	Ile	Pro	Lys	Glu	Leu	Lys	Lys	His	Met	Gln	Ala		
305					310					315					

What is claimed is:

1. A method for the production of isoprenoid compounds comprising: contacting a transformed host cell under suitable growth conditions with an effective amount of a carbon source whereby an isoprenoid compound is produced, said transformed host cell comprising a nucleic acid molecules encoding SEQ ID NOs: 2 under the control of suitable regulatory sequences.

2. A method according to claim 1 wherein the transformed host cell is selected from the group consisting of *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Rhodococcus*, *Streptomyces*, *Escherichia*, *Pseudomonas*, *Methylomonas*, *Methylobacter*, *Alcaligenes*, *Synechocystis*, *Anabaena*, *Thiobacillus*, *Methanobacterium* and *Klebsiella*.

3. A method according to claim 1 wherein said transformed host cell is a methanotrophic bacteria.

4. A method according to claim 3 wherein said methanotrophic bacteria:

- (a) grows on a C1 carbon substrate selected from the group consisting of methane and methanol; and
- (b) comprises a functional Embden-Meyerof carbon pathway, said pathway comprising a gene encoding a pyrophosphate dependent phosphofructokinase enzyme.

10

5. A method according to claim 3 wherein the methanotrophic bacteria is selected from the group consisting of *Methylomonas*, *Methylobacter* and *Methanobacterium* and the carbon source is selected from the group consisting of methane and methanol.

15

6. A method according to claim 4 wherein said methanotrophic bacteria is *methylomonas* 16a ATCC PTA 2402.

20

7. A method according to claim 1 wherein the transformed host cell is selected from the group consisting of soybean, rapeseed, sunflower, cotton, corn, tobacco, alfalfa, wheat, barley, oats, sorghum, rice, *Arabidopsis*, cruciferous vegetables, melons, carrots, celery, parsley, tomatoes, potatoes, strawberries, peanuts, grapes, grass seed crops, sugar beets, sugar cane, beans, peas, rye, flax, hardwood trees, softwood trees, and forage grasses.

25

8. A method according to claim 1 wherein the carbon source is selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, carbon dioxide, methanol, methane, formaldehyde, formate, and carbon-containing amines.

30

* * * * *

not claimed

-continued

385	390	395	400
His Glu Asp Glu Thr	Gly Tyr Pro Asp	Asp Leu Leu Ala	Glu Asp Gly
	405	410	415
Leu Ser Val Ser Asp	Asn Ser Pro Ile	Ala Pro Gly Glu	Thr Arg Thr
	420	425	430
Val Glu Val Thr Ala	Ser Asp Ala Ala	Trp Glu Val Tyr	Arg Leu Ala
	435	440	445
Asp Leu Ile Tyr Asp	Pro Asp Ser Arg	Phe Ala Gly Leu	Met Phe Phe
	450	455	460
Trp Asp Glu Asn Gly	Asn Arg Gln Met	Thr Met Val Asp	Ala Pro Leu
	465	470	475
Ile Pro Thr Phe Ile			480
	485		

<210> SEQ ID NO 81
 <211> LENGTH: 1429
 <212> TYPE: DNA
 <213> ORGANISM: METHYLOMONAS SP.
 <400> SEQUENCE: 81

```

cggtagtgcct aacacatgca agtgcgaacgc tgaaggggtgc ttgcacctgg atgagtggcg 60
gacgggtgag taatgcatag gaatctgcct attagtgggg gataacgtgg ggaactcac 120
gctaataccg catacgtctc acggaggaaa gccggggacc ttcgggctcg gcgctaatag 180
atgagcctat gtcggattag ctagtgtgtg gggtaaaggc ctaccaaggc gacgatccgt 240
agctggtctg agaggatgat cagccacact gggactgaga caccgcccag actcctacgg 300
gaggcagcag tggggaatat tggacaatgg gcgcaagcct gatccagcaa taccgctgtg 360
gtgaagaagg cctgaggggtt gtaaagcact ttcaatggga aggaacacct atcggttaat 420
accggtaga ctgacattac ccatacaaga agcaccggct aactccgtgc cagcagcccg 480
ggtaatacgg aggggtgcaag cgtaaatcgg aattactggg cgtaaacgtg gcgtaggcgg 540
ttttttaagt cagatgtgaa agccctgggg ttaacctggg aactgcattt gatactgggg 600
aactagagtt gagtagagga gagggtgaatt tcaggtgttag cgggtgaaatg cgtagagatc 660
tgaaaggaaa ccagtggcga agcggtctct ctggactcaa actgacgtg aggtacgaaa 720
gcgtgggtag caaacaggat tagataccct ggtagtccac gccgtaaaacg atgtcaacta 780
accgttgggt tcttaaagaa cttagtgggt gagctaactg attagttga cgcctgggg 840
agtacggccg caaggctaaa actcaaatga attgacgggg gcccgacaaa gcgggtggagc 900
atgtgggtta attcgatgca acgcgaagaa ccttacctac ccttgacatc ctcggaactt 960
gtcagagatg acttgggtgc ttccgggaacc gagagacagg tgctgcattg ctgtcgtcag 1020
ctcgtgtcgt gagatgttgg gtttaagtcct gtaacgagcg caaccttat ccttagttgc 1080
cagcgcgta tggcggaac tctagggaga ctgccgtgta taacccggag gaaggtgggg 1140
acgacgtcaa gtcacatggt cccttatggg tagggctaca cacgtgctac aatggtcggt 1200
acagaggggt gcgaactcgc gagagccagc caatcccaaa aagccgatcc tagtccggat 1260
tgacgtctgc aactcgactt gcatgaagtc ggaatcgcta gtaatcgcg atcagaatgc 1320
cgcggtgaat acgttcccg gccttgata caccgccgt cacaccatgg gagtgggtg 1380
caaaagaagt aggtagtta accttcggga gggcgcttac cactttgtg 1429

```

not claimed

What is claimed is:

1. A pure isolate of a high growth methanotrophic bacterial strain which:

- grows on a C1 carbon substrate selected from the group consisting of methane and methanol; and
- comprises a functional Embden-Meyerhof carbon pathway, said pathway comprising a gene encoding a pyrophosphate dependent phosphofructokinase enzyme, the gene comprising an isolated nucleic acid molecule encoding the amino acid sequence as set forth in SEQ ID NO: 6.

2. A high growth methanotrophic bacterial strain according to claim 1 wherein the strain contains a functional Entner-Doudoroff carbon pathway.

3. A bacterial strain according to claim 1 having at least one gene encoding a fructose biphosphate aldolase enzyme.

4. A bacterial strain according to claim 3 wherein at least one gene encodes a fructose biphosphate aldolase enzyme having the amino acid sequence selected from the group consisting of SEQ ID NO:16 and SEQ ID NO:18.

5. The bacterial strain of claim 1 wherein the strain is a *Methylomonas* sp.

6. The bacterial strain of claim 5 having a 16s RNA profile as set forth in SEQ ID NO:81.

7. The bacterial strain of claim 1 wherein, when the C1 carbon substrate is methanol, the strain produces glycogen comprising at least about 50% dry weight of biomass.

8. The bacterial strain of either claim 1 or claim 7 wherein the methanol concentration in the medium is about 2.5% (vol/vol).

9. The bacterial strain of any of claims 1 or 2 having a yield of greater than 1.0 grams of cell mass per gram of methane consumed.

10. The bacterial strain of any of claims 1 or 2 having a yield of greater than 0.5 grams of cell mass per gram of methane consumed.

11. The bacterial strain of any of claims 1 or 2 having a carbon conversion efficiency of greater than 40 g/mol methane/g/mol biomass.

12. The bacterial strain of any of claims 1 or 2 having a carbon conversion efficiency of greater than 64 g/mol methane/g/mol biomass.

13. A pure isolate of a high growth methanotrophic bacterial strain which grows on a C1 carbon substrate selected from the group consisting of methanol and methane, comprising the 16s RNA sequence as set forth in SEQ ID NO:81 and having at least one gene encoding a pyrophosphate dependent Phosphofructokinase enzyme.

14. A pure isolate of a high growth methanotrophic bacterial strain having the ATCC designation PTA 2402.

* * * * *